recombination and B cell development (Galloway et al., 2016**). Thus, by comparison to pre-B cells the mechanisms and genes that control proliferation in pro-B cells remain poorly understood.**

Transcription factors act to determine which genes are transcribed and the tempo of transcription. A significant body of work has identified a network of transcription factors that control the development and identity of B cells (Busslinger, 2004**). Amongst these, FOXO1 is essential for progression after the pro-B cell stage and to induce expression of *Rag* genes (**Dengler et al., 2008**;** Amin and Schlissel, 2008**). After transcription, numerous RBPs (**Gerstberger et al., 2014**) control messenger RNA (mRNA) expression and coordinate functionally related genes into mRNA regulons (**Keene, 2007**). These post-transcriptionally controlled networks are more challenging to identify because they**

A

Ctrl
CD79a$^{+/+}$;
Ptbp1$^{fl/fl}$;Ptbp2$^{fl/fl}$

P1sKO
CD79a$^{Cre/+}$;
Ptbp1$^{fl/fl}$;Ptbp2$^{+/+}$

P1P2dKO
CD79a$^{Cre/+}$;
Ptbp1$^{fl/fl}$;Ptbp2$^{fl/fl}$

Rag2$^{-/-}$

B



A



cells (Figure 4A**) since** *Ptbp2* **transcripts in P1sKO pro-B cells included exon 10 and were no longer**

mRNA abundance and AS patterns in P1P2dKO pro-B cells (Figure 4—source datas 1 **and** 2**). E2A (encoded by** *Tcf3*) **and IKAROS are two transcription factors with AS (**Schjerven et al., 2013**;** Sun and Baltimore, 1991**) important for B cell ontogeny. In human, PTBP1 regulates AS of** *Tcf3* **(**Yamazaki et al., 2019**). We found similar AS patterns of** *Ikaros* **and** *Tcf3* **in P1P2dKO compared to control pro-B cells (**Figure 4—source data 2**).** *Ebf1* **mRNA abundance was reduced ~1.5 fold in P1P2dKO compared to control and P1sKO pro-B cells (**Figure 5A**), but this should not impact B cell development because** *Ebf1* **haploinsufficiency results in normal B cell ontogeny (**Vilagos et al., 2012**;** György et al., 2012**).** *Foxo1* **mRNA abundance was reduced ~1.7 fold in P1P2dKO compared to P1sKO and control pro-B cells (**Figure 5A**). PTBP1 bound directly to the 3'UTR of** *Foxo1* **(**Figure 5B**), indicating a direct role of PTBP1 stabilising** *Foxo1* **mRNA.** *Foxo1* **deficient pro-B cells do not develop further and have reduced IL-7 receptor expression (**Dengler et al., 2008**). Indeed,** *Il7r* **mRNA abundance was reduced ~1.4 fold in P1P2dKO compared to P1sKO and control pro-B cells (**Figure 5A**). However, IL-7 receptor staining was similar between P1P2dKO and control pro-B cells (**Figure 5C**), ruling out a role for a reduction of IL-7 receptor in the developmental defect of P1P2dKO pro-B cells. FOXO1 promotes RAG expression (**Amin and Schlissel, 2008**) but** *Rag1* **and** *Rag2* **mRNA abundance and AS were unaffected in P1P2dKO pro-B cells (**Figure 4—figure supplement 2E **and** Figure 4—source data 2**). These data indicate that the known essential elements of the B**


Il7r
0.002
0.008
16
12
8
4
0
x10³

(Figure 6—figure supplement 1C**), which are indicative of an increased cell size and are consistent with a high biosynthetic capacity and a predisposition to enter S-phase.**

To understand if the increased proportions of P1P2dKO pro-B cells in S-phase were due to an

5'UTR (Supplementary file 2**) although** *Cdkn1b* **mRNA is expressed in mitogen-activated B cells. Thus, additional layers of regulation are expected to contribute to the dynamic regulation of p27 protein abundance across the different phases of the cell cycle observed in the absence of PTBP1 and PTBP2.**



proportions of p-RB and p-SAMDH1-positive cells were reduced in P1P2dKO mice compared to control and P1sKO mice (Figure 7C and D **and** Figure 7—figure supplement 1B**). This indicates a requirement for PTBP1 to achieve sufficient CDK1 activity to enter mitosis. Thus, PTBP1 is essential for controlling the activity of CDKs in different phases of the cell cycle in pro-B cells, to limit entry into S-phase and to promote entry into mitosis.**

B on ro s n K pr ss on o ph s nr g ors n pro B s

GO enrichment analysis of genes differentially expressed in P1P2dKO compared to control and P1sKO pro-B cells identified an enrichment for ribosome biogenesis and CDK inhibitors but it did not detect further terms directly related with proliferation. Therefore to identify PTBP1 targets directly implicated in the progression through different phases of the cell cycle we assessed changes in mRNA abundance of genes known to be highly expressed specifically in the S and G2/M-phases (Giotti et al., 2019**) between FrB pro-B cells of the different genotypes. S- and G2/M-associated transcripts were not globally increased in P1P2dKO FrB pro-B cells compared to control and P1sKO FrB pro-B cells (**Figure 8A **and** Figure 8—source data 1**). This indicates that the majority of the changes observed in the transcriptome of P1P2dKO compared to control and P1sKO pro-B cells are not the result of comparing populations with different proliferative status, but are reflective of an altered pro-B cell transcriptome. Thus, instead of gene sets, it was possible that individual genes with rate limiting properties for cell cycle progression, which could also control the expression of CDK-inhibitors and CDK activities, were directly regulated by PTBP1. Therefore, we considered further genes affected in the absence of PTBP1 and PTBP2 but not in the absence of PTBP1 alone that are known to regulate the cell cycle.**

We identified four genes associated with S-phase entry with direct PTBP1 binding and changes at the levels of mRNA abundance and AS. Two genes (*Myc* **and** *Ccnd2***) whose proteins promote entry into S-phase, showed increased mRNA abundance in P1P2dKO compared to P1sKO and control FrB pro-B cells (**Figure 8B**). Two further genes (***Btg2* **and** *Rbl1***), whose proteins inhibit S-phase entry, had reduced mRNA abundance in P1P2dKO compared to control and P1sKO FrB pro-B cells (**Figure 8B**). PTBP1 bound to the 3'UTR of** *Myc* **(**Figure 8—figure supplement 1**), and we confirmed an increase**

Log2 Fold Change

cells (Figure 9A**). CDC25B is a phosphatase promoting G2/M transition by activating CDK1 (**Boutros et al., 2007**).** *Cdc25b* **AS was unaffected by PTBP1 and PTBP2 absence but PTBP1 bound to the** *Cdc25b* **3'UTR (**Figure 4—source data 2 **and** Figure 8—figure supplement 1**). Thus, PTBP1 probably promotes** *Cdc25b* **expression by binding to its 3'UTR and enhancing its stability in pro-B cells.**

There were three genes (*Ect2***,** *Kif2c* **and** *Kif22***) whose changes in mRNA abundance could be**

The molecular mechanisms of PTBP action on the cell cycle mRNA regulon that we assessed were limited to the regulation of AS and mRNA abundance. PTBPs are known, in addition, to control polyadenylation site choice and IRES-mediated translation of certain transcripts (Hu et al., 2018**). Therefore, we can expect that additional roles for PTBP1 regulating translation efficiencies and polyadenylation site usage in pro-B cells may emerge. Moreover, by suppressing the inclusion of non-conserved cryptic exons that generate NMD targets (**Ling et al., 2016**;** Sibley et al., 2016**), PTBPs may play a broader role in maintaining the global 'fidelity' of the transcriptome. Cryptic**

deoxyuridine, cat# B5002-500mg, Sigma) labelling experiments mice were injected first with 1 mg EdU (cat #E10415, ThermoFisher Scientific) intraperitoneally. One hour later, the same mice were injected with 2 mg BrdU and the mice were killed 1 hr and 40 min after the injection with EdU.

Fo ~ o r

Single cell suspensions were prepared from spleens and lymph nodes by passing the organs through cell strainers with 70 μm and 40 μm pore sizes in cold RPMI-1640 (cat# R8758, Sigma) with 2% fetal

PTBP1 iCLIP was carried out previously from mitogen-activated B cells (splenic B cells stimulated with LPS for 48 hr) (Monzón-Casanova et al., 2018**). The data generated were re-analysed to map reads using a splicing-aware software. Reads from five PTBP1 iCLIP libraries were mapped to the GRCm38.p5 mouse genome from Gencode with STAR (v 2.5.4b) (**Dobin et al., 2013**). Reads were de-duplicated using random barcodes included in the library preparation and xlink-sites (**Supplementary file 1**) and clusters of binding (**Supplementary file 2**) were identified with iCount https://icount.readthedocs.io/en/latest/cite.html as previously described (**Monzón-Casanova et al., 2018**). Detection of a binding site with iCLIP (**König et al., 2010**) is highly dependent on the abundance of the RNA, therefore all replicates were pooled together to identify xlink sites and clusters of binding. Xlink sites and clusters of PTBP1 binding were assigned to transcripts and genomic features with the following hierarchy: CDS, 3'UTR, 5'UTR, intron, ncRNA using the Mus_musculus. GRCm38.93.gtf annotation from Ensembl (**Supplementary files 1 **and** 2**). mRNA abundance analysis mRNAseq libraries were trimmed with Trim Galore (v 1.15 https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with default parameters and mapped with Hisat2 (v 2.1.0) (**Kim et al., 2015**) with -p 7 t –**phred33-quals –no-mixed –no-discordant **parameters, the Mus_musculus. GRCm38 genome and known splice sites from the Mus_musculus.GRCm38.90 annotation. Read counts mapped over genes were counted with HTSeq (v0.10.0) (**Anders et al., 2015**) with f bam -r name -s no -a 10 parameters and the Mus_musculus.GRCm38.93.gtf annotation from Ensembl. Reads mapping to immunoglobulin genes (including V, D, J gene segments, light and heavy immunoglobulin genes and T cell receptor genes) were excluded before DESeq2 analysis. Differences in mRNA abundance were computed with DESeq2 (v1.22.1) (**Love et al., 2014**) by extracting differences between different genotypes in pair-wise comparisons using 'apeglm' method as a shrinkage estimator (**Zhu et al., 2019**). Information on the sex of the mice from which the mRNAseq libraries were generated was included in the design formula in addition to the genotype (design = ~**

biological replicates mapping to either the included or skipped alternative splicing event in at least one of the two conditions analysed. Only alternative splicing events with the highest inclusion level difference were kept out of AS events which had the same genomic coordinates in the AS event (Figure 4—source data 2**). rMATS considers five different types of alternative splicing events: skipped exons (SE), mutually exclusive exons (MXE), alternative 5' and 3' splice sites (A5SS and A3SS, respectively) and retained introns (RI) (**Figure 4—figure supplement 2C**). Proportions of differential alternative splicing events were defined as bound by PTBP1 (**Figure 4—figure supplement 2D **and** Figure 4—source data 2**) with different criteria depending on the type of AS event.**

An SE was bound by PTBP1 if PTBP1 clusters (Supplementary file 2**) were found on the SE, 500 nucleotides upstream or downstream of the AS SE, on either of the constitutive flanking exons or in 500 nucleotides downstream or upstream of the upstream and downstream constitutive exons, respectively. An A5SS was bound if PTBP1 clusters were found on the longest exon containing the A5SS, on the downstream intronic 500 nucleotides of the A5SS, on the downstream constitutive flanking exon or the 500 intronic nucleotides upstream of the downstream constitutive flanking exon. An A3SS was bound by PTBP1 if PTBP1 clusters were found on the longest exon containing the A3SS, the 500 intronic nucleotides upstream of the A3SS, the upstream constitutive flanking exon or the 500 intronic nucleotides downstream of the upstream constitutive flanking exon. An MXE was bound by PTBP1 if PTBP1 clusters were found on either MXE, the upstream or downstream 500 intronic nucleotides of either MXE, the upstream or downstream constitutive flanking exons or the downstream or upstream 500 intronic nucleotides from the upstream or downstream constitutive flanking exons, respectively. A RI was bound by PTBP1 if PTBP1 clusters were found on the RI or on either constitutive flanking exon.**

Events with changes in AS in P1P2dKO FrB pro-B cells compared to control and P1sKO FrB pro-B cells (Figure 4E**) were identified by hierarchical clustering using complete linkage clustering of the Euclidean distances between Z-scores of each AS event calculated from the rMATS inclusion levels for the 15 mRNAseq libraries (five biological replicates per genotype). AS events found in any of the three pairwise comparisons (**Figure 4B**) were used in the hierarchical clustering.**

Gene ontology analysis

Genes belonging to different clusters based on differences in mRNA abundance or AS patterns between the P1P2dKO FrB pro B cells and the other genotypes (P1sKO and Controls) (Figure 4D and E **and** Figure 5—source data 1

Ethics

Animal experimentation: Mice were bred and maintained in the Babraham Institute Biological Support Unit. Since the opening of this barrier facility (2009), no primary pathogens or additional agents listed in the FELASA recommendations have been confirmed during health monitoring surveys of the stock holding rooms. Ambient temperature was ~19-21˚C and relative humidity 52%. Lighting was provided on a 12-hour light: 12-hour dark cycle including 15 min 'dawn' and 'dusk' periods of subdued lighting. After weaning, mice were transferred to individually ventilated cages with 1-5 mice per cage. Mice were fed CRM (P) VP diet (Special Diet Services) ad libitum and received seeds (e.g. sunflower, millet) at the time of cage-cleaning as part of their environmental enrichment. All mouse experimentation was approved by the Babraham Institute Animal Welfare and Ethical Review Body (UK Home Office Project Licence /P4D4AF812). Animal husbandry and experimentation complied with existing European Union and United Kingdom Home Office legislation and local standards.

Decision letter and Author response

Decision letter https //doi.org/10.7 /eLife. 3 7 .sa1
Author response https //doi.org/10.7 /eLife. 3 7 .sa2

A on s

Supple entary files

- **Supplementary file 1. PTBP1 binding sites (xlinks).**
- **Supplementary file 2. PTBP1 binding sites (clusters).**
- **Supplementary file 3. Key resources table.**
- **Transparent reporting form**

Data availability

E, Cunningham AF, Gonzalez-Porta M, Andrews SR, Bunik VI, Zarnack K, Curk T, Kontoyiannis DL, Ule J, Turner M				
Diaz-Muñoz MD, Bell SE, Fairfax K, Monzon-Casanova E, Cunningham AF, Gonzalez-Porta M, Andrews SR, Bunik VI, Zarnack K, Curk T, Kontoyiannis DL, Ule J, Turner M	201	WT_LPS2	https //www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM1 2011	NCBI Gene Expression Omnibus, GSM1 2011
Diaz-Muñoz MD, Bell SE, Fairfax K, Monzon-Casanova E, Cunningham AF, Gonzalez-Porta M, Andrews SR, Bunik VI, Zarnack K, Curk T, Kontoyiannis DL, Ule J, Turner M	201	WT_LPS3	https //www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM1 2011	NCBI Gene Expression Omnibus, GSM1 2011
Diaz-Muñoz MD, Bell SE, Fairfax K, Monzon-Casanova E, Cunningham AF, Gonzalez-Porta M, Andrews SR, Bunik VI, Zarnack K, Curk T, Kontoyiannis DL, Ule J, Turner M	201	HuR- dependent regulation of mRNA splicing is essential for the B cell antibody response	https //www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE 212	NCBI Gene Expression Omnibus, GSE 212
Ling JP, Chhabra R, Merran JD, Schaughency PM, Wheelan SJ, Corden JL, Wong PC	201	PTBP1 and PTBP2 Repress Nonconserved Cryptic Exons	https //www.ncbi.nlm.nih.gov/bioproject/PRJNA30 7 32	NCBI BioProject, PRJNA30 7 32

Cr r A

or , Matus-Nicodemos R, Valentín-Acevedo A, Covey LR. 2016. The RNA-Binding protein, polypyrimidine Tract-Binding protein 1 (PTBP1) is a key regulator of CD4 T cell activation. *PLOS ONE* :e0158708. DOI: https://doi.org/10.1371/journal.pone.0158708, PMID: 27513449

Z, Dordai DI, Lee J, Desiderio S. 1996. A conserved degradation signal regulates RAG-2 accumulation during cell division and links V(D)J recombination to the cell cycle. *Immunity* :575–589. DOI: https://doi.org/10.1016/S1074-7613(00)80272-1, PMID: 8986717

, Zheng S, Han A, Lin C-H, Stoilov P, Fu X-D, Black DL. 2014. The splicing regulator PTBP2 controls a program of embryonic splicing required for neuronal maturation. *eLife* :e01201. DOI: https://doi.org/10.7554/eLife.01201

, Chhabra R, Merran JD, Schaughency PM, Wheelan SJ, Corden JL, Wong PC. 2016. PTBP1 and PTBP2 repress nonconserved cryptic exons. *Cell Reports* :104–113. DOI: https://doi.org/10.1016/j.celrep.2016.08.071, PMID: 27681424

or n , Schwartz S, Clark TA, Hollander D, Tan LY, Spellman R, Gordon A, Schweitzer AC, de la Grange P, Ast G, Smith CW. 2010. Position-dependent alternative splicing activity revealed by global profiling of alternative splicing events regulated by PTB. *Nature Structural & Molecular Biology* :1114–1123. DOI: https://doi.org/10.1038/nsmb.1881, PMID: 20711188

, Wendling O, Masjkur J, Jäger M, Münster C, Anastassiadis K, Stewart AF, Solimena M. 2011. PTBP1 is required for embryonic development before gastrulation. *PLOS ONE* :e16992. DOI: https://doi.org/10.1371/journal.pone.0016992, PMID: 21423341

n X , Baltimore D. 1991. An inhibitory domain of E12 transcription factor prevents DNA binding in E12 homodimers but not in E12 heterodimers. *Cell* :459–470. DOI: https://doi.org/10.1016/0092-8674(91)90653-G, PMID: 1846322

s E, Nakayama K, Nakayama KI. 2007. Cyclin D2 translocates p27 out of the nucleus and promotes its degradation at the G0-G1 transition. *Molecular and Cellular Biology* :4626–4640. DOI: https://doi.org/10.1128/MCB.00862-06, PMID: 17452458

n Y, Whitfield P, Llorian M, Monzon-Casanova E, Diaz-Munoz MD, Turner M, Smith CW. 2015. Generation of functionally distinct isoforms of PTBP3 by alternative splicing and translation initiation. *Nucleic Acids Research* :5586–5600. DOI: https://doi.org/10.1093/nar/gkv429, PMID: 25940628

o pson B , Bhansali R, Diebold L, Cook DE, Stolzenburg L, Casagrande AS, Besson T, Leblond B, Désiré L, Malinge S, Crispino JD. 2015. DYRK1A controls the transition from proliferation to quiescence during lymphoid development by destabilizing cyclin D3. *The Journal of Experimental Medicine* :953–970. DOI: https://doi.org/10.1084/jem.20150002, PMID: 26008897

r pn C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and cufflinks. *Nature Protocols* :562–578. DOI: https://doi.org/10.1038/nprot.2012.016

r n .